

**Mixing and Feeding Directions  
Medicated Feeds Containing 0.0384%  
Dichlorvos**

Prepare a complete feed containing 0.0384% dichlorvos (174 mg/lb) according to the following mixing chart.

Pounds of Complete Feed	Pounds of ATGARD C to be Added to Feed
500	2
1000	4
2000	8

See package insert for complete use directions.

**For Swine up to 70 lbs Body Weight:**

Feed as the sole ration for 2 consecutive days. Use the following table as a guide for estimating consumption of the medicated feed:

Weight of Pigs (lbs)	Amount of Medicated Feed to be Fed in 2 Days (lbs per Pig)
30	4.0
40	5.5
50	6.5
60	7.5
70	8.4

**For swine from 70 lbs to market weight:** Feed as the sole ration at the rate of 8.4 lbs of feed per head until all the medicated feed has been consumed.

**For boars, open or bred gilts or sows:** Feed as the sole ration at the rate of 4.2 lbs of feed per head per day for 2 consecutive days. In all cases, after the allotted medicated feed has been consumed, resume feeding regular rations.

There is no pre-slaughter withdrawal period when used at the recommended dosage level.

**Warning:** Keep out of the reach of children. Avoid contact with skin. ATGARD C is a cholinesterase inhibitor. If human or animal poisoning should occur, immediately consult a physician or a veterinarian. Atropine is antidotal.

Medicated feeds containing ATGARD C must bear the following statements:

**Note** - Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

**Warning**

1. Do not use in animals other than swine.
2. Do not mix in feeds to be pelleted nor with pelleted feed. (Required only on medicated feed supplements.)
3. Do not allow fowl access to this feed or to feces from treated animals.
4. Do not soak feed or feed as wet mash. Feed must be fed dry.
5. Dichlorvos is a cholinesterase inhibitor. Do not use this product in animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals. If human or animal poisoning should occur immediately consult a physician or veterinarian. Atropine is antidotal.
6. Do not reuse any of the containers or container materials used in packaging (product). All packaging materials must be destroyed after (product) has been used. (Required only on medicated feed supplements.)

TO OPEN CARTON SLIT ALONG THIS SEAM

TO OPEN CARTON SLIT ALONG THIS SEAM

**To Deworm Swine**

For the removal and control of mature, immature and/or fourth stage larvae of the whipworm (*Trichuris suis*), nodular worm (*Oesophagostomum* sp.), large roundworm (*Ascaris suum*), and thick stomach worm (*Ascaraps strongylina*) occurring in the gastrointestinal tracts of pigs, boars, open or bred gilts and sows.

**Mixing and Feeding Directions**

ATGARD C can be mixed in meal or crumble feeds after crumbles have been manufactured. Do not mix in feeds to be pelleted nor with pelleted feed.

**Individual Rations**

Each package of ATGARD C (dichlorvos) Swine Wormer contains a plastic scoop. A level scoop holds 15.2 grams ATGARD C (1460 mg dichlorvos), the recommended dewormer dosage for swine over 70 lbs.

To deworm **one** open or bred gilt or sow, boar, or pig from 70 lbs to market weight, thoroughly mix **one** level scoop of ATGARD C into the usual daily ration of each animal. After the allotted medicated feed has been consumed, resume feeding regular rations.

This package contains sufficient ATGARD C (dichlorvos) Swine Wormer to individually deworm 239 sows.



Lot No.:

Exp. Date:

Boehringer Ingelheim Vetmedica, Inc.  
St. Joseph, MO 64506 U.S.A.

**610101L-01-9801**  
Code 610151

Deworming Mixing Directions – Complete Feeds			
Pounds Complete Feed	Dichlorvos		Pounds Atgard® C to be Added Per Ton of Feed
	mg/lb	%	
2000	174	0.0384	8.0
2000	239	0.0528	11.0

To Improve Litter Production Efficiency Mixing Directions – Complete Feeds			
Pounds Complete Feed per Head Daily	Dichlorvos in Feed		Pounds Atgard® C to be Added Per Ton of Feed
	mg/lb	%	
4	250	0.0552	11.5
5	200	0.0442	9.2
6	167	0.0368	7.7

**Boehringer Ingelheim Vetmedica, Inc.**  
2621 North Belt Highway • St. Joseph, MO 64506-2002  
Information 800-325-9167 • Orders 800-325-9167 • Fax 816-236-2717  
www.bi-vetmedica.com