

# Cephalovac® VEWT

**Encephalomyelitis Vaccine  
Eastern, Western, and Venezuelan, Killed Virus, Tetanus Toxoid**

## Description

Encephalomyelitis Vaccine Eastern, Western, and Venezuelan, Killed Virus, Tetanus Toxoid. Cephalovac® VEWT contains gentamicin and thimerosal as preservatives.

## Features

Provides broad spectrum active immunity against Eastern, Western, and Venezuelan Equine Encephalomyelitis and tetanus.

Contains killed viruses, tetanus toxoid and a potent, surface-active adjuvant.

Produced in stable cell-line cultures.

## Benefits

Safe to use on all healthy horses.

Adjuvant stimulates antigen processing.

## Advantages

Cephalovac VEWT is a killed virus vaccine that protects against Eastern, Western and Venezuelan Equine Encephalomyelitis and tetanus.

## Dosage & Administration

Using aseptic technique inject 2 mL IM. For primary vaccination administer a second 2 mL dose in 4 to 6 weeks using a different injection site. Revaccinate annually and prior to anticipated exposure, using a single 2 mL dose. Use a separate, sterile 1½ inch needle for each injection.

## Precautions

Store out of direct sunlight at 35° - 45° F (2° - 7° C). Shake well. Protect from freezing. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur following use. Antidote: Epinephrine.

## Uses

Cephalovac VEWT is recommended for vaccination of all types and ages of healthy horses against Eastern, Western, and Venezuelan Equine Encephalomyelitis and tetanus.

## Warning

This product has been tested under laboratory conditions and shown to meet all Federal standards for safety and efficacy in normal healthy animals. This level of performance may be affected by conditions such as stress, weather, nutrition, disease, parasitism, other treatments, individual idiosyncrasies or impaired immunological competency. These factors should be considered by the user when evaluating product performance or freedom from reactions.

## Product Information

Name	Code	Size
Cephalovac® VEWT	131-615	2 mL 1 dose
	131-625	20 mL 10 doses

## Highlights

- Sold only to licensed veterinarians.



Serial No.:  
Exp. Date:

**Indications:** For the vaccination of all types and ages of healthy horses against Eastern, Western and Venezuelan Equine Encephalomyelitis and tetanus.

**Administration and Dosage:** Shake well before using. Inject 2 mL deep into muscle. Revaccination required. See carton for full directions.

**Caution:** Store at 35-45°F (2-7°C). Protect from freezing. Use entire contents when first opened. Contains gentamicin and thimerosal as

## Encephalomyelitis Vaccine,

Eastern, Western, and Venezuelan, Killed Virus

## Tetanus Toxoid

**Cephalovac® VEWT**

10 Doses/20 mL

preservatives. Do not vaccinate within 21 days before slaughter. Anaphylactoid reactions may occur following use.

**Antidote:** Epinephrine.

**Boehringer Ingelheim Vetmedica, Inc.**

**St. Joseph, MO 64506 U.S.A.**

**U.S. Veterinary License No. 124**

Code 131-325; 31303-02



**Boehringer  
Ingelheim**

**Boehringer Ingelheim Vetmedica, Inc.**

**2621 North Belt Highway • St. Joseph, MO 64506-2002**

**Information 800-325-9167 • Sales Service 800-325-9167 • Fax 816-236-2717**

**[www.bi-vetmedica.com](http://www.bi-vetmedica.com)**