

Serial No.:  
Exp. Date:

**Indications:** Recommended for the vaccination of healthy, susceptible gilts and sows as an aid in the reduction of disease associated with Porcine Reproductive and Respiratory Syndrome Virus, reproductive form. The duration of immunity is at least 4 months post vaccination or throughout gestation.

**Directions:** Rehydrate the vaccine by adding the accompanying liquid diluent to the vaccine vial. Shake well and use immediately. See insert for complete directions.

**Dosage:** Using aseptic technique, administer a single 5 mL dose intramuscularly to gilts and sows 4-6 weeks prior to breeding. Revaccinate prior to each subsequent breeding.

**Precautions:** Store out of direct sunlight at a temperature between 35-45°F (2-

## Porcine Reproductive and Respiratory Syndrome Vaccine

Reproductive Form,  
Modified Live Virus

**ReproCyc® PRRS**

5 Doses/25 mL

7°C). Use entire contents when first opened. Do not vaccinate within 21 days of slaughter. Do not vaccinate pregnant gilts and sows or boars. Burn containers and all unused contents. Anaphylactoid reactions may occur. **Antidote:** Administer Epinephrine. Contains Neomycin as a preservative.

**Boehringer Ingelheim  
Vetmedica, Inc.**  
St. Joseph, MO 64506  
(800) 325-9167  
U.S. Veterinary License No. 124  
424702N-00



**Boehringer  
Ingelheim**

Serial No.:  
Exp. Date:

**Indications:** Recommended for the vaccination of healthy gilts and sows as an aid in the prevention and control of disease caused by porcine parvovirus, *Erysipelothrix rhusiopathiae*, and *Leptospira canicola*, *grippotyphosa*, *hardjo*, *icterohaemorrhagiae* and *pomona*.

**Directions:** Shake well. Administer 5mL intramuscularly to gilts and sows 4-6 weeks prior to breeding. Repeat in 3-4 weeks. Revaccinate with a single dose prior to each subsequent breeding.

**Precautions:** Store out of direct sunlight at 35-45°F (2-7°C). Do not freeze. Use entire contents when

**Parvovirus Vaccine  
Erysipelothrix  
Rhusiopathiae-Leptospira  
Canicola-Grippotyphosa-  
Hardjo-Icterohaemorrhagiae  
Pomona Bacterin, Killed Virus**

5 Doses/25 mL

first opened. If anaphylactoid reaction occurs, epinephrine is symptomatic treatment. Do not vaccinate within 21 days of slaughter. Contains amphotericin B, penicillin, streptomycin and thimerosal as preservatives.

For Veterinary Use Only.

**Boehringer Ingelheim  
Vetmedica, Inc.**  
St. Joseph, MO 64506  
(800) 325-9167  
U.S. Veterinary License No. 124  
472802N-00



**Boehringer  
Ingelheim**

**Boehringer Ingelheim Vetmedica, Inc.**

2621 North Belt Highway • St. Joseph, MO 64506-2002

Information 800-821-7467 • Customer Service 800-325-9167 • Fax 816-236-2717

Website [www.bi-vetmedica.com](http://www.bi-vetmedica.com)