

ReproCyc® PRRS-PLE

PRRS, Porcine Parvovirus, *Leptospira* and Erysipelas Vaccine

Description

ReproCyc® PRRS-PLE is a convenient prebreeding combination vaccine composed of a modified live PRRS component and a nonvirucidal, inactivated Parvo *Leptospira* and *Erysipelothrix rhusiopathiae* diluent component.

Features

Greatly reduces reproductive failure due to PRRS virus.

Reduces PRRS viremia in challenged animals.

Aids in the prevention and control of disease caused by Porcine parvovirus, *Erysipelothrix rhusiopathiae* and 5 strains of *Leptospira*.

Proven cross-protection against homologous and heterologous (Lelystad virus) PRRS challenge.

Benefits

Save time and labor vs. non-combination products.

Proven cross-protection.

Convenient combination fits gilt introduction vaccination protocols, as well as ongoing sow vaccination programs.

Advantages

Easy to administer pre-breeding combination.

PRRS cross-protection proven by challenge.

Dosage & Administration

ReproCyc PRRS-PLE is a vaccine administered as a convenient 5 mL dose to gilts and sows 4-6 weeks prior to breeding. Revaccinate with a parvovirus/*Leptospira*/erysipelas vaccine 3-4 weeks following initial vaccination. Revaccinate with a single dose of ReproCyc PRRS-PLE vaccine prior to subsequent breedings and only vaccinate gilts and sows in PRRS-positive herds.

Precautions

Store out of direct sunlight at 35°- 45° F (2° - 7° C). Do not freeze. Use entire contents when first opened. If anaphylactoid reaction occurs epinephrine is symptomatic treatment.

Uses

ReproCyc PRRS-PLE is designed specifically for sows and gilts. It is recommended for the vaccination of healthy sows and gilts as an aid in the reduction of disease caused by porcine reproductive and respiratory syndrome virus reproductive form, porcine parvovirus, the five major types of leptospirosis affecting breeding-age swine including *Leptospira canicola*, *L. grippotyphosa*, *L. hardjo*, *L. icterohaemorrhagiae* and *L. pomona* and *Erysipelothrix rhusiopathiae*.

Warning

Do not vaccinate within 21 days of slaughter.

Do not vaccinate pregnant sows or gilts, or boars of breeding age. For use only in PRRS virus positive herds as directed.

Highlights

- Convenient, cost-efficient.
- Proven cross-protection against homologous and heterologous (Lelystad Virus) PRRS challenge.
- Safe, proven and effective.

Product Information

Name	Code	Size
ReproCyc® PRRS-PLE	173-711	5 doses 20 mL
	173-721	20 doses 100 mL



Serial No.:
Exp. Date:

Indications: Recommended for the vaccination of healthy, susceptible gilts and sows as an aid in the reduction of disease associated with Porcine Reproductive and Respiratory Syndrome Virus, reproductive form. The duration of immunity is at least 4 months post vaccination or throughout gestation.

Directions: Rehydrate the vaccine by adding the accompanying liquid diluent to the vaccine vial. Shake well and use immediately. See insert for complete directions.

Dosage: Using aseptic technique, administer a single 5 mL dose intramuscularly to gilts and sows 4-6 weeks prior to breeding. Revaccinate prior to each subsequent breeding.

Precautions: Store out of direct sunlight at a temperature between 35-45°F (2-

Porcine Reproductive and Respiratory Syndrome Vaccine

Reproductive Form,
Modified Live Virus

ReproCyc® PRRS

5 Doses/25 mL

7°C). Use entire contents when first opened. Do not vaccinate within 21 days of slaughter. Do not vaccinate pregnant gilts and sows or boars. Burn containers and all unused contents. Anaphylactoid reactions may occur. **Antidote:** Administer Epinephrine. Contains Neomycin as a preservative.

**Boehringer Ingelheim
Vetmedica, Inc.
St. Joseph, MO 64506
(800) 325-9167
U.S. Veterinary License No. 124
424702N-00**



**Boehringer
Ingelheim**

Serial No.:
Exp. Date:

Indications: Recommended for the vaccination of healthy gilts and sows as an aid in the prevention and control of disease caused by porcine parvovirus, *Erysipelothrix rhusiopathiae*, and *Leptospira canicola*, *grippotyphosa*, *hardjo*, *icterohaemorrhagiae* and *pomona*.

Directions: Shake well. Administer 5mL intramuscularly to gilts and sows 4-6 weeks prior to breeding. Repeat in 3-4 weeks. Revaccinate with a single dose prior to each subsequent breeding.

Precautions: Store out of direct sunlight at 35-45°F (2-7°C). Do not freeze. Use entire contents when

Parvovirus Vaccine Erysipelothrix Rhusiopathiae-Leptospira Canicola-Grippotyphosa- Hardjo-Icterohaemorrhagiae Pomona Bacterin, Killed Virus

5 Doses/25 mL

first opened. If anaphylactoid reaction occurs, epinephrine is symptomatic treatment. Do not vaccinate within 21 days of slaughter. Contains amphotericin B, penicillin, streptomycin and thimerosal as preservatives.

For Veterinary Use Only.

**Boehringer Ingelheim
Vetmedica, Inc.
St. Joseph, MO 64506
(800) 325-9167
U.S. Veterinary License No. 124
472802N-00**



**Boehringer
Ingelheim**

Boehringer Ingelheim Vetmedica, Inc.

2621 North Belt Highway • St. Joseph, MO 64506-2002

Information 800-821-7467 • Customer Service 800-325-9167 • Fax 816-236-2717

Website www.bi-vetmedica.com