

Non-steroidal anti-inflammatory drug for use in dogs and cats only

**Warning:** Not for use in humans. Keep this and all medications out of reach of children. Refer to package insert for additional information.

Store at controlled room temperature, 68–77°F (20–25°C).

Manufactured by:

**Boehringer Ingelheim Vetmedica, Inc.**  
St. Joseph, MO 64506 U.S.A.



601304L-01-0410  
Code 601311

**Metacam**<sup>®</sup>

(meloxicam)

**5 mg/mL Solution for Injection**

**Caution:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**Net Contents: 10 mL**

**Indications: Dogs:** For the control of pain and inflammation associated with osteoarthritis.

**Cats:** For the control of postoperative pain and inflammation associated with orthopedic surgery, ovariohysterectomy and castration when administered prior to surgery.

Refer to package insert for complete dosage and administration information.

Lot No.:

Exp. Date:



**Boehringer  
Ingelheim**

## Package Insert for Cats

NADA 141-219, Approved by FDA

# Metacam® (meloxicam)


## 5 mg/mL Solution for Injection

Non-steroidal anti-inflammatory drug for use in dogs and cats only

**Caution:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

### Description:

Meloxicam is a non-steroidal anti-inflammatory drug (NSAID) of the oxicam class. Each mL of this sterile product for injection contains meloxicam 5.0 mg, alcohol 15%, glycofurool 10%, poloxamer 188 5%, sodium chloride 0.6%, glycine 0.5% and meglumine 0.3%, in water for injection, pH adjusted with sodium hydroxide and hydrochloric acid.



### Indications:

**Cats:** For the control of postoperative pain and inflammation associated with orthopedic surgery, ovariohysterectomy and castration when administered prior to surgery.

### Dosage and Administration:

**Cats:** Administer a **single, one-time subcutaneous dose** of Metacam 5 mg/mL Solution for Injection to cats at a dose of 0.14 mg/lb (0.3 mg/kg) body weight. **Use of additional meloxicam or other NSAIDs is contraindicated. (See Contraindications).** To ensure accuracy of dosing, the use of a 1 mL graduated syringe is recommended.

### Contraindications:

Cats with known hypersensitivity to meloxicam and other NSAIDs should not receive Metacam 5 mg/mL Solution for Injection. Additional doses of meloxicam or other NSAIDs in cats are contraindicated, as no safe dosage for repeated NSAID administration has been established (See Animal Safety).

**Warnings:** Not for use in humans. Keep this and all medications out of reach of children. Consult a physician in case of accidental ingestion by humans. For subcutaneous (SQ) injectable use in cats. Do not use IV in cats.

▶ **Do not administer a second dose of meloxicam.** ◀

▶ **Do not follow meloxicam dose with any other NSAID.** ◀

When administering any NSAID, appropriate laboratory testing to establish hematological and serum biochemical baseline data is recommended prior to use in dogs and cats. All cats should undergo a thorough history and physical examination before administering meloxicam. Do not repeat dose in cats.

**Owner should be advised to observe their cats for signs of potential drug toxicity.**

### Precautions:

The safe use of Metacam 5 mg/mL Solution for Injection in cats younger than 4 months of age, cats used for breeding, or in pregnant or lactating queens has not been evaluated. Meloxicam is not recommended for use in cats with bleeding disorders, as safety has not been established in cats with these disorders. Safety has not been established for intravenous (IV) or intramuscular (IM) use in cats. When administering Metacam 5 mg/mL Solution for Injection, use a syringe of appropriate size to ensure precise dosing. As a

class, cyclo-oxygenase inhibitory NSAIDs may be associated with gastrointestinal, renal and hepatic toxicity. Sensitivity to drug-associated adverse events varies with the individual patient. Patients at greatest risk for renal toxicity are those that are dehydrated, on concomitant diuretic therapy, or those with existing renal, cardiovascular, and/or hepatic dysfunction. Concurrent administration of potentially nephrotoxic drugs should be carefully approached. Anesthetic drugs may affect renal perfusion; approach concomitant use of anesthetics and NSAIDs cautiously. NSAIDs may inhibit the prostaglandins that maintain normal homeostatic function. Such anti-prostaglandin effects may result in clinically significant disease in patients with underlying or preexisting disease that has not been previously diagnosed.

In cats, appropriate monitoring procedures should be employed during all surgical procedures. The use of parenteral fluids during surgery is recommended to decrease potential renal complications when using NSAIDs. If additional pain medication is needed after the single dose of meloxicam, a non-NSAID class of analgesic may be necessary. In one study<sup>1</sup>, one cat in each NSAID treatment group had increased intraoperative hemorrhage.

Since many NSAIDs possess the potential to produce gastrointestinal ulceration, concomitant use of Metacam 5 mg/mL Solution for Injection with other antiinflammatory drugs, such as corticosteroids, should be avoided or closely monitored in cats. Consider appropriate washout times when switching from corticosteroid use to meloxicam in cats. **As a single use product in cats, meloxicam should not be followed by additional NSAIDs or corticosteroids.** The use of concomitantly protein-bound drugs with Metacam 5 mg/mL Solution for Injection has not been studied in cats. Commonly used protein-bound drugs include cardiac, anticonvulsant and behavioral medications. The influence of concomitant drugs that may inhibit metabolism of Metacam 5 mg/mL Solution for Injection has not been evaluated. Drug compatibility should be monitored in patients requiring adjunctive therapy. The effect of cyclo-oxygenase inhibition and the potential for thromboembolic occurrence or a hypercoagulable state has not been studied.

### Adverse Reactions:

**Cats:** A field study involving 138 cats was conducted. Of the 72 cats receiving Metacam 5 mg/mL Solution for Injection, six cats (8.3%) experienced post-treatment elevated serum blood urea nitrogen (BUN) levels. The pre-treatment values were in the normal range. Of the 66 cats in the butorphanol treatment group, no cats experienced post-treatment elevated serum blood urea nitrogen levels. Nine cats (12.5%) receiving Metacam 5 mg/mL Solution for Injection had post-treatment anemia. Pre-treatment, these cats all had hematocrit and hemoglobin values in the normal range. Four cats (6.1%) in the butorphanol treatment group had post-treatment anemia. All but one cat, who had a mild anemia pre-treatment (hematocrit=21% and hemoglobin=7.0 g/dL) had normal pretreatment values. Twenty-four hours after the injection with Metacam 5 mg/mL Solution for Injection, one cat experienced pain upon palpation of the injection site.

### Foreign Experience:

**Repeated use in cats has been associated with acute renal failure and death.** In studies used for the foreign approval of Metacam 5 mg/mL Solution for Injection in cats, lethargy, vomiting, inappetance, and transient pain immediately after injection were noted.

Diarrhea and fecal occult blood have also been reported.

To report suspected adverse reactions, to obtain a Material Safety Data Sheet, or for technical assistance, call 1-866-METACAM (1-866-638-2226).

**Information For Cat Owners:** Meloxicam, like other NSAIDs, is not free from adverse reactions. Owners should be advised of the potential for adverse reactions and be informed of the clinical signs associated with NSAID intolerance. Adverse reactions may include vomiting, diarrhea, lethargy, decreased appetite and behavioral changes. Cat owners should be advised when their pet has received a meloxicam injection. Cat owners should contact their veterinarian immediately if possible adverse reactions are observed.

**Clinical Pharmacology:** Meloxicam has nearly 100% bioavailability after subcutaneous injection in cats. The terminal elimination half life after a single dose is estimated to be approximately 15 hrs (+/-10%) in cats.

Peak drug concentrations of 1.1 mcg/mL can be expected to occur within 1.5 hours following a 0.3 mg/kg subcutaneous injection in cats. The volume of distribution (Vd $\lambda$ ) in cats is approximately 0.27 L/kg, with an estimated total systemic clearance of 0.013 L/hr/kg. The drug is 97% bound to feline plasma proteins.

**Effectiveness:**

**Cats:** The effectiveness of Metacam 5 mg/mL Solution for Injection was demonstrated in a masked field study involving a total of 138 cats representing various breeds. This study used butorphanol as an active control. Cats received either a single subcutaneous injection of 0.3 mg/kg Metacam 5 mg/mL Solution for Injection or 0.4 mg/kg butorphanol prior to onychectomy, either alone or in conjunction with surgical neutering. All cats were premedicated with acepromazine, induced with propofol and maintained on isoflurane. Pain assessment variables evaluated by veterinarians included additional pain intervention therapy, gait/lameness score, analgesia score, sedation score, general impression score, recovery score, and visual analog scale score. Additionally, a cumulative pain score, which was the summation of the analgesia, sedation, heart rate and respiratory rate scores was evaluated. A palpometer was used to quantify the pain threshold.

A substantial number of cats required additional intervention in the 0-24 hour postsurgical period, with the majority of these interventions taking place within the first hour. Therefore, the percentage of cats in each group that received one or more interventions was designated as the primary assessment variable. Approximately half of the cats in each group received a pain intervention as a result of the first (time 0) post-surgical evaluation, i.e., extubation. At this point, the need to provide a pain intervention was not statistically significant between the two groups (p=0.7215). However, the median number of interventions was one per cat in the meloxicam group and two per cat in the butorphanol group and this difference was statistically significant (p=0.0021). The statistical evaluation supports the conclusion that the meloxicam test article is noninferior to the butorphanol active control. Forty-eight of the 72 cats in the meloxicam group received one or more interventions (66.7%), and 47 of 66 cats in the butorphanol group received one or more interventions (71.2%). The number of interventions administered to the meloxicam group was less than the butorphanol group at 1, 3, 5, 8, 12, and 24 hours post-surgery.

Cats receiving Metacam 5 mg/mL Solution for Injection showed improvement in the pain assessment variables.

**Animal Safety:**

**Cats: 3 Day Target Animal Safety Study** - In a three day safety study, subcutaneous Metacam 5 mg/mL Solution for Injection administration to healthy cats at up to 1.5 mg/kg (5X the recommended dose) resulted in vomiting in three cats (1 of 6 control cats and 2 of 6 cats in 5X) and loose stools in four cats (2 of 6 control cats and 2 of 6 cats in 5X).

Fecal occult blood was detected in ten of the twenty four cats, including two cats in the control group. This was not a dose-related event. Clinically significant hematologic changes seen included increased PT and APTT in two cats (1 of 6 control cats and 1 of 6 cats in 5X), and elevated white blood cell counts in cats having renal or GI tract lesions. Serum chemistry changes observed included decreased total protein in four of 24 cats (1 of 6 cats in 1X, 2 of 6 cats in 3X and 1 of 6 cats in 5X), concomitant increases in blood urea nitrogen (BUN) and creatinine values in 2 of 6 cats in 5X.

Histological examination revealed gastrointestinal lesions ranging from inflammatory cell infiltration of the mucosa of the GI tract to erosions. Mesenteric lymphadenopathy was identified in 1 of 6 cats in 1X. Renal changes ranged from dilated medullary (2 of 6 cats in 1X, 1 of 6 cats in 3X, and 1 of 6 cats in 5X) and cortical (3 of 6 cats in 1X, 1 of 6 cats in 3X, and 3 of 6 cats in 5X) tubules and inflammation (2 of 6 cats in 1X, 2 of 6 cats in 3X, and 2 of 6 cats in 5X) or fibrosis (2 of 6 cats in 3X and 2 of 6 cats in 5X) of the interstitium to necrosis of the tip of the papilla (5 of 6 cats in 5X).

**Subsequent oral dosing** - In a nine day study with three treatment groups, Metacam 5 mg/mL Solution for Injection was given as a single subcutaneous injection using doses of 0 mg/kg (saline injection), 0.3 mg/kg and 0.6 mg/kg on Day 0. Metacam Oral Suspension, 1.5 mg/mL or saline was then administered orally once-daily at the same respective dose (0.3 or 0.6 mg/kg) for eight consecutive days. Clinical adverse reactions included vomiting, diarrhea, lethargy, and decreased food consumption in the treated groups, and one day of diarrhea in one control cat. The gross necropsy report includes observation of reddened GI mucosa in 3 of 4 cats in the 0.3 mg/kg group and 1 of 4 cats in the 0.6 mg/kg group. All saline-treated cats were normal. By Day 9, one cat in both the 0.3 mg/kg group and the 0.6 mg/kg group died and another cat in the 0.3 mg/kg group was moribund. The cause of death for these cats could not be determined, although the pathologist reported pyloric/duodenal ulceration in the cats in 0.6 mg/kg group. The safety studies demonstrate a narrow margin of safety.

**Injection Site Tolerance** - Histopathology of the injection sites revealed hemorrhage and inflammation, myofiber atrophy, panniculitis, fibrin deposition, and fibroblast proliferation. These findings were present in cats in all groups, with the 3X cats having the most present. No safe repeat dose has been established in cats.

**Storage Information:** Store at controlled room temperature, 68-77°F (20-25°C).

**How Supplied:**

Metacam 5 mg/mL Solution for Injection: 10 mL vial

**Reference:**

1 Slingsby L.S., A.E. Waterman-Pearson. Comparison between meloxicam and carprofen for postoperative analgesia after feline ovariohysterectomy. *Journal of Small Animal Pract* (2002)43:286-289.

**Manufactured by:**

Boehringer Ingelheim Vetmedica, Inc.  
St. Joseph, MO 64506 U.S.A.

Metacam® is a registered trademark of Boehringer Ingelheim Vetmedica GmbH, licensed to Boehringer Ingelheim Vetmedica, Inc.

601307L-01-0410  
Code 601311  
Revised 10-2004